

Policy: MP185

Section: Medical Benefit Policy

Subject: Chemosensitivity and Chemoresistance Assays

I. Policy: Chemosensitivity and Chemoresistance Assays

II. Purpose/Objective:

To provide a policy of coverage regarding Chemosensitivity and Chemoresistance Assays

III. Responsibility:

- A. Medical Directors
- B. Medical Management

IV. Required Definitions

1. Attachment – a supporting document that is developed and maintained by the policy writer or department requiring/authoring the policy.
2. Exhibit – a supporting document developed and maintained in a department other than the department requiring/authoring the policy.
3. Devised – the date the policy was implemented.
4. Revised – the date of every revision to the policy, including typographical and grammatical changes.
5. Reviewed – the date documenting the annual review if the policy has no revisions necessary.

V. Additional Definitions

Medical Necessity or Medically Necessary means Covered Services rendered by a Health Care Provider that the Plan determines are:

- a. appropriate for the symptoms and diagnosis or treatment of the Member's condition, illness, disease or injury;
- b. provided for the diagnosis, and the direct care and treatment of the Member's condition, illness disease or injury;
- c. in accordance with current standards of good medical treatment practiced by the general medical community.
- d. not primarily for the convenience of the Member, or the Member's Health Care Provider; and
- e. the most appropriate source or level of service that can safely be provided to the Member. When applied to hospitalization, this further means that the Member requires acute care as an inpatient due to the nature of the services rendered or the Member's condition, and the Member cannot receive safe or adequate care as an outpatient.

Medicaid Business Segment

Medical Necessity shall mean a service or benefit that is compensable under the Medical Assistance Program and if it meets any one of the following standards:

- (i) The service or benefit will, or is reasonably expected to, prevent the onset of an illness, condition or disability.
- (ii) The service or benefit will, or is reasonably expected to, reduce or ameliorate the physical, mental or development effects of an illness, condition, injury or disability.
- (iii) The service or benefit will assist the Member to achieve or maintain maximum functional

capacity in performing daily activities, taking into account both the functional capacity of the Member and those functional capacities that are appropriate for members of the same age.

DESCRIPTION: A chemosensitivity assay refers to any in vitro laboratory analysis that is performed especially to determine whether tumor growth is inhibited by a known chemotherapeutic agent or a panel of drugs. Conversely, chemoresistance assays are used to determine specific chemotherapeutic agents to which a patient's tumor will not adequately respond. The use of in vitro chemotherapy sensitivity and resistance assays is thought to aid the physician in choosing the most appropriate and successful treatment for their patients.

EXCLUSIONS: The Plan does **NOT** provide coverage for Chemosensitivity and Chemoresistance Assays (including, but not limited to, the ChemoFx® assay) for any indication because it is considered **experimental, investigational and unproven**. The Geisinger Technology Assessment Committee evaluated this technology and concluded that there is insufficient evidence in the peer-reviewed published medical literature to establish the effectiveness of this test on health outcomes when compared to established tests or technologies.

Note: A complete description of the process by which a given technology or service is evaluated and determined to be experimental, investigational or unproven is outlined in **MP 15 - Experimental Investigational or Unproven Services or Treatment**.

CODING ASSOCIATED WITH:

The following codes are included below for informational purposes and may not be all inclusive. Inclusion of a procedure or device code(s) does not constitute or imply coverage nor does it imply or guarantee provider reimbursement. Coverage is determined by the member specific benefit plan document and any applicable laws regarding coverage of specific services. Please note that per Medicare coverage rules, only specific CPT/HCPCS Codes may be covered for the Medicare Business Segment. Please consult the CMS website at www.cms.gov or the local Medicare Administrative Carrier (MAC) for more information on Medicare coverage and coding requirements

81535 Oncology (gynecologic), live tumor cell culture and chemotherapeutic response by DAPI stain and morphology, predictive algorithm reported as a drug response score; first single drug or drug combination

81536 Oncology (gynecologic), live tumor cell culture and chemotherapeutic response by DAPI stain and morphology, predictive algorithm reported as a drug response score; each additional single drug or drug combination ChemoFX

Current Procedural Terminology (CPT®) © American Medical Association: Chicago, IL

LINE OF BUSINESS:

Eligibility and contract specific benefits, limitations and/or exclusions will apply. Coverage statements found in the line of business specific benefit document will supersede this policy. For Medicare, applicable LCD's and NCD's will supercede this policy. For PA Medicaid Business segment, this policy applies as written.

REFERENCES:

Geisinger Technology Assessment Triage Committee Review, Chemotherapy Sensitivity & Resistance Assays June, 2006.

Geisinger Technology Assessment Committee Review, Chemotherapy Sensitivity & Resistance Assays July, 2009.

Link KH, Kornmann M, Butzer U, Ieder G, Sunelaitis E, Pillasch J, Salonga D, Danenberg KD, Danenberg PV, Beger HG. Thymidylate Syntheses Quantitation and In Vitro Chemosensitivity testing predicts responses and survival of patients with isolated nonresectable liver tumor receiving hepatic arterial infusion chemotherapy. Cancer 2000;89:288-96.

Blue Cross and Blue Shield Association Technology Evaluation Center: Chemotherapy sensitivity and resistance assays: TECT Assesments 17:1-39, 2002.

Schrag D, Garewal HS, Burstein HJ, Samson DJ, Von Hoff DD, Somerfield MR. American Society of Clinical Oncology Technology Assessment: Chemotherapy Sensitivity and Resistance Assays. J Clin Oncol 22:3631-3638.

Samson DJ, Seidenfeld J, Ziegler K, Aronson N. Chemotherapy sensitivity and resistance assays: A systematic review. J Clin Oncol 22:3618-3630.

Winifred S. Hayes. Hayes Directory (online). In vitro extreme drug resistance assays in cancer treatment. Hayes Inc. Lansdale Pa. Current as of March 28, 2005.

Winifred S. Hayes. Hayes Directory (online). In vitro chemosensitivity assays in cancer treatment. Hayes Inc. Lansdale Pa. Current as of February 7, 2006

Loizzi V, Chan JK, Osann K, Cappuccini F, DiSaia PJ, Berman ML. Survival Outcomes in patients with recurrent ovarian cancer who were treated with chemoresistance assay-guided chemotherapy. *Am J Obstet Gynecol* 2003;189:1307-7.

Sharma S, Neale MH, Nicolantonio FD, Knight LA, Whitehouse PA, Mercer SJ, Higgins BR, Lamont A, Osborne R, Hindley AC, Kurbacher CM, Cree IA. Outcome of ATP-based tumor chemosensitivity assay directed chemotherapy in heavily pre-treated recurrent ovarian carcinoma. *BMC Cancer* 2003;3:19.

Maenpaa JU, Heinonen E, Hinkka SM, Karnani P, Klemi PJ, Korpijaakko TA, Kuoppala TA, Laine MA, Nuoranne EK et al. The subrenal capsule assay in selecting chemotherapy for ovarian cancer: a prospective randomized trial. *Gynecol Oncol*. 1995 Jun;57(3):294-8.

Staib P, Staltmeier E, Neurohr K, Cornely O, Reiser M, Schinkothe T. Prediction of individual response to chemotherapy in patients with acute myeloid leukaemia using the chemosensitivity index C₁. *British Journal of Haematology* 2004;128:783-791.

Eltabbakh GH. Extreme drug resistance assay and response to chemotherapy in patients with primary peritoneal carcinoma. *J. Surg. Oncol.* 2000;73:148-152.

d'Amato TA, Landreneau RJ, McKenna RJ, Santos RS, Parker RJ. Prevalence of in vitro extreme chemotherapy resistance in resected nonsmall-cell lung cancer. *Ann Thorac Surg* 2006;81:440-7.

Cortazar P and Johnson BE. Review of the efficacy of individualized chemotherapy selected by invitro drug sensitivity testing for patients with cancer. *J Clin Oncol* 1999;17:1625-1631.

Cloven NG, Kyshtoobayeva A, Burger RA, Yu IR, Fruehauf JP. In vitro chemoresistance and biomarker profiles are unique for histologic subtypes of epithelial ovarian cancer. *Gynecol Oncol*. 2004 Jan;92 (1):160-6.

Des Guetz G, Uzzan B, Nicolas P, et al. Microsatellite instability does not predict the efficacy of chemotherapy in metastatic colorectal cancer. A systematic review and meta-analysis. *Anticancer Res*. 2009;29(5):1615-1620.

Julian C. Schink, MD, Larry J. Copeland, MD Point: Chemosensitivity Assays Have a Role in the Management of Recurrent Ovarian Cancer. *J Natl Compr Canc Netw* 2011;9:115-120

LCD L36634 In Vitro Chemosensitivity & Chemoresistance Assays. <https://www.novitas-solutions.com/policy/mac-ab/l32571.html>

Tian C, Sargent DJ, Krivak TC, et al. "Evaluation of a chemoresponse assay as a predictive marker in the treatment of recurrent ovarian cancer: further analysis of a prospective study ." *British Journal of Cancer* 2014

Krivak TC, Lele S, Richard S, et al. A chemoresponse assay for prediction of platinum resistance in primary ovarian cancer. *Am J Obstet Gynecol*. 2014;211(1):68.e1-e8

ECRI Institute. Hotline Response. In Vitro Chemosensitivity Assays for Predicting Ovarian Cancer Response to Chemotherapy, July 2014

This policy will be revised as necessary and reviewed no less than annually.

Devised: 07/2006

Revised: 04/11, 10/12 (Local Intermediary Non-Coverage position); 8/15 (clarification)

Reviewed: 12/07, 07/09 (TAC review), 7/10, 8/11, 8/12, 8/13, 8/14, 7/16, 7/17, 7/18